

Unveiling the Complexity of Red Blood Cells: Insights into Structure, Properties and Functions

Alexandra AVRAM¹, Aurora MOCANU¹, Gheorghe TOMOAIA^{2,3},
Cristina Lavinia ROȘOIU^{1,4}, Cristina Teodora DOBROTĂ^{1,4},
Diana Alexandra FLOREA¹, Maria TOMOAIA-COTISEL^{1,3}

¹ Babes-Bolyai University of Cluj-Napoca, Faculty of Chemistry and Chemical Engineering,
Research Centre of Physical Chemistry, 11 Arany Janos Str., RO-400028, Cluj-Napoca,
Romania

² Iuliu Hatieganu University of Medicine and Pharmacy, Department of Orthopedics and
Traumatology, 47 General Traian Mosoiu Str., RO-400132, Cluj-Napoca, Romania

³ Academy of Romanian Scientists, 3 Ilfov Str., RO-050044, Bucharest, Romania

⁴ Babes-Bolyai University of Cluj-Napoca, Faculty of Biology and Geology, Department
of Molecular Biology and Biotechnology, 44 Republicii Str., RO-400015, Cluj-Napoca,
Romania

*Corresponding author: Maria Tomoaia-Cotisel, mcotisel@gmail.com
maria.tomoaia@ubbcluj.ro

Abstract. *Considering the basic function of red blood cells (RBC, erythrocytes) as carriers of oxygen and carbon dioxide throughout the bloodstream, as well as their possible secondary activities, RBCs deserve more consideration. The current work attempts to serve as a summary of RBC properties, both well-established and less well-established, with a focus on pathologies and drug interactions. This review is especially important given the recent trend of employing erythrocytes as vehicles for targeted medication delivery.*

Keywords: red blood cells, functions, pathology, interactions with drugs

DOI [10.56082/annalsarscibio.2023.1.129](https://doi.org/10.56082/annalsarscibio.2023.1.129)

1. Introduction

Human blood is formed by plasma and a variety of cells, namely white (*leukocytes*: neutrophils, lymphocytes – B and T cells, monocytes, basophils, eosinophils and macrophages), red (*erythrocytes*) and platelets (*thrombocytes*) [1]. Although all blood components are quite complex, with new information being continuously discovered, they all have their basic functions. While less than 1% of the entire blood content, leukocytes aid in fighting against infections and disease [2]. Platelets, on the other hand, help stop bleeding by interacting with blood

clotting proteins [3]. Last, but not least, red blood cells (RBCs) are the predominant cells within blood, making up to 40-45% of its volume, being quite simple in structure and possessing no internal organelles [4]. For a better understanding, Figure 1 presents a graphic representation of blood composition.




Figure 1. Blood composition

Blood components can interact with one another and are extremely dynamic. Erythrocytes, for example, can influence the movement of white blood cells by excluding them from the bulk solution [5]. It was demonstrated that the shape of erythrocytes affects their ability (through organized aggregation) to induce the radial migration of white blood cells. RBCs also interact with platelets, leading to the latter's concentration being higher towards the wall of the blood vessel. This type of interaction can be either of a turning (platelets do not make direct RBC contact and turn around in the direction they came from) or a crossing type (platelets roll over erythrocytes and proceed with their way, leading to a mild RBC deformation) [6]. Macrophages are deeply connected with red blood cells, being involved in both erythropoiesis (development and differentiation of RBCs

in the bone marrow) and erythrophagocytosis (removal of defective RBCs) [7]. As it would be impossible to convey, in a concise manner, all in depth aspects regarding every single blood component, specific functions and interactions, this review will focus on conveying some aspects regarding red blood cells.

2. The basics on red blood cells

The basic function of RBC is the transportation of oxygen from the lungs towards tissues as well as carbon dioxide back to the lungs while travelling around 300 km within the bloodstream [8]. This is achieved through the hemoglobin (Hb) which is a tetramer comprised of four polypeptide globin chains. The three types of hemoglobin (HbA, HbA2, HbF) are derived from the four types of globin chains, namely α , β , γ , δ , with each subunit containing an iron atom bound to a heme molecule [9].

The transportation of gases is due to the iron, with each hemoglobin having the ability to transport about four molecules of either carbon dioxide or oxygen. Most carbon dioxide, however, is transported as bicarbonate ions (HCO_3^-), where the H_2CO_3 deprotonation in water is a result of the CO_2 and H_2O reaction, which is mediated by carbonic anhydrase [10]. Nonetheless, RBCs confront some obstacles even though their role is quite straightforward. These biconcave disks with a thickness and diameter of around 2-2.5 μm and 7-8 μm , respectively, must be able to squeeze through capillaries as small as 3 μm in diameter, as well as through slits of less than 1 μm in width in the reticuloendothelial system [11]. Their high plasticity is a crucial parameter to ensure that oxygen is properly delivered. Any disturbance in the shape-maintaining ability of RBCs would lead to hemolytic anemia. The deformation property of red blood cells is of course in part due to their unique biconcave shape, but also the composition and organization of the cell membrane (52% protein, 40% lipids, 8% carbohydrates), namely cytoskeleton and lipid bilayer [12, 13]. There are many intricacies concerning the structure of the erythrocyte membrane, but briefly, the lipid bilayer is composed of phospholipids, namely phosphatidylethanolamine and phosphatidylserine (inner layer) and phosphatidylcholine and sphingomyelin (outer layer) with the intervening space being occupied by cholesterol [14]. On the other hand, spectrin, ankyrin, protein 4.1R, actin, as well as actin associated protein form the RBC cytoskeleton which binds to the bilayer through protein complexes [15]. Given that the precise position of several components is still unknown [15], Figure 2 provides a simplified schematic representation of these aspects.




Figure 2. Simplified schematic representation of the red blood cell membrane composition

More details regarding the red blood cell membrane components in Figure 2 and their specific functions are presented in Table 1.

Table 1. Details regarding the major components of RBC membrane

Type	Component	Role and details	Ref
<i>Proteins</i>	Band 3 (AE1/ SLC4A1/ capnophorin)	- transmembrane protein anchoring the membrane skeleton - Anion transporter - the most abundant protein in membrane consisting of 911 amino acids	[16,17]
	Glycophorins A, B	- are encoded by 2 genes that are similar (GYPA, GYPB) - carry the MN and Ss blood groups - glycophorin A is the most abundant	[18,19]
	Glycophorins C, D	- are encoded by a similar gene (GYPC) - glycophorin C carries the Gerbich blood group system - glycophorin C aids in maintaining the shape of RBC - glycophorin C is specific to humans - glycophorin D is conserved for all apes	[18,20]
	Spectrin α and β	- cytoskeletal proteins maintaining the shape and flexibility of the membrane - spectrin can affect rheological properties of blood - there are 2 α and 5 β spectrin subunits - α and β spectrin subunits mainly associate through hydrophobic bonds supported by electrostatic forces	[21-23]
	Actin	- short actin filaments (approx. 37 nm in length) connect with spectrin tetramers forming RBC skeleton - control the mechanics of RBC	[24-26]

	Ankyrin	<ul style="list-style-type: none"> - cytoskeletal protein - anchors the cytoskeleton to the RBC membrane - attaches band 3 and protein 4.2 to β spectrin and skeleton - Ankyrin's compression into a spring-like state may be a factor in the distinctive shape and flexibility of RBC cells. 	[27,28]
	4.2	<ul style="list-style-type: none"> - peripheral membrane protein - 5% of membrane content - its absence leads to hereditary spherocytosis - binds to the N-terminal cytoplasmic domain of band 3 - interacts with ankyrin 	[29,30]
	4.1	<ul style="list-style-type: none"> - forms a ternary complex with spectrin and actin - the attachment to glycophorin C bridges the protein network to the bilayer 	[31]
<i>Lipids</i>	Phospholipids	<ul style="list-style-type: none"> - arranged in an asymmetrical manner into a bilayer with a thickness of two molecules - extracellular surface is more abundant in phosphatidylcholine with sphingomyelins and a small amount of phosphatidylserine - the inner part of the bilayer is mostly abundant in amino phospholipids, primarily consisting of phosphatidylethanolamines, phosphatidylserines and phosphatidylcholine with smaller levels of sphingomyelins - aid with the fluidity of RBC 	[32,33]
	Neutral lipids	<ul style="list-style-type: none"> - comprised of mostly cholesterol that is intercalated in between the phospholipid molecules - aid with the fluidity of RBC 	[32]

As mentioned previously, carbohydrates form about 8% of the dry mass of the erythrocyte membrane. They are mostly relevant for researchers as part of glycoproteins, glycopeptides or glycolipids. The carb residues bound to glycophorins for example can be better observed in Figure 2. Within the red blood cell membrane, carbohydrates are present in many structures such as the Lewis, MN and ABO blood-group specific antigens [34]. The ABO blood grouping, for example, is conditioned by differences in glycoproteins (protein carbohydrate complexes). Type A blood has type A glycoprotein, type B blood has type B glycoproteins, type AB is characterized by the presence of both and type O by the presence of none.

Again, while red blood cells are very complex and many intricacies will not be discussed here, aquaporins must be mentioned. They were first discovered in human RBCs [35-39] and later purified [40] and further studied in relation to their water transport property [41, 42]. Aquaporins or water channel proteins have a primary function of transporting water across cell membranes as a response to

osmotic gradients. The presence of these proteins is integral for RBCs and their proper functioning.

2.1. Transfusions and further prospects

Erythrocytes have a life span of around 120 days before being removed by the spleen. This dramatically changes however when it comes to shelf life. Blood in blood banks, typically stored at 4 °C, is viable for up to 42 days. Of course, during storage some cells might undergo hemolysis at a swifter pace than others, releasing hemoglobin (Hb) in the bag or once they have undergone transfusion [43]. Transfusion of RBCs at the end of their shelf life is unadvisable as it can lead to an increased risk in mortality [44]. This poses a conundrum for patients suffering from trauma-related severe blood loss or different types of blood conditions requiring periodic transfusions. This is a particular concern when it comes to O negative blood types. Cryopreservation would be one solution to such problems.

Erythrocytes preserved in 40-percent (weight/volume) glycerol are approved for storage at -80 °C up to 10 years by the USA-based Federal Drug Administration (FDA) [45]. According to Chang et al. the FDA consents to the employment of such RBCs (processed in ACP 215 Automated Cell Processor (Haemonetics Corp., Baintree, MA, USA)) for only up to 2 weeks since the removal of glycerol [46]. According to a double-blind randomized trial [47], cryopreserved red blood cells (CPRBC) are superior to liquid preserved ones (LPRBC), leading to the absence of an inflammatory response, having an attenuated fibrinolytic state as well as an increased level of 2,3-DPG. Here, the 2,3-DPG (2,3-diphosphoglycerate) is a crucial parameter that influences the proficiency by which hemoglobin releases oxygen to the tissues.

However, while hemolysis is an unwanted factor when it comes to preserving human life, it can have alternative uses. The low shelf life and degradation of RBCs outside of the human body can be used in forensic science towards determining the time at which a certain criminal act was committed. The characteristics of erythrocytes undergo degradation from the moment they encounter the environmental conditions outside of the human body. Hemoglobin for example, is present in 2 forms within the body (with and without oxygen) and while outside of the body it becomes completely saturated with oxygen, auto-oxidizes to methemoglobin, and then suffers a denaturation process to hemichrome [48].

The autoxidation of hemoglobin was reported to comprise an initial fast decay followed by a slower one [49]. One study [50] researched the use of a non-destructive method, namely Attenuated Total Reflection – Fourier Transform Infrared (ATR-

FTIR) spectroscopy, for bloodstain identification in mimicked crime scenarios. This however proved more applicable for older stains, particularly 7-85 days.

The physical characteristics, namely morphology and elasticity, can also be analyzed in comparison to those of the standard, fresh erythrocyte, to determine the time at which a certain crime had taken place. Numerous analysis techniques have been employed to determine the age of a certain bloodstain. One such technology is hyperspectral imaging (HSI), which is favored because it is less invasive, although it is not without drawbacks. Edelman et al. [51] simulated a crime scene with blood stains of various ages (0.1; 2; 15; 40 and 200 days) and measured the temporal behavior of certain blood fractions such as oxyhemoglobin (HbO₂), methemoglobin (MetHb) and hemichrome (HC). This study reports that a mean difference between the actual age of the stain and the measured one is placed in the 0.9-6-day range.

This discrepancy has been extensively explored in the forensic scientific literature, and various analysis approaches have been tested. High resolution electron microscopies might offer an insight into the visible morphological characteristics of red blood cells. Nonetheless, this might not always lead to enough information. However, Atomic Force Microscopy (AFM) can provide an advantage as it not only allows for morphological studies through simple imaging, but it can also give insight into the viscoelastic properties of those RBCs through force measurements. One specific study [52] observed the changes in erythrocytes over a span of 4 weeks by Atomic Force Microscopy. Here, while morphology was not necessarily altered, the elasticity parameters of RBC did decrease over time, a change attributed to drying and coagulation. Another study [53] working with blood from donors (4 females and 4 males; 20-40 years old) managed to identify statistically quantifiable changes in some morphological characteristics of erythrocytes, namely cell perimeter, height, area, and volume during a 28-day experiment.

A decrease in elasticity after 21 days could probably be attributed to an alteration of lipid conformation within the membrane. It was determined that the substrate used for sample deposition is an important factor in these types of time-dependent analyses. An additional information to consider is that the indentation process is a crucial aspect of accurately determining RBCs Young Modulus parameters by AFM. As reported by Smijs et al. [54] the material from which the used cantilever is made does influence the measurements, especially as the stiffness of erythrocytes increases over time.

3. Erythrocyte abnormalities in terms of morphology

Erythrocytes are formed from stem cells found in bone marrow. To reiterate, they appear as biconcave disks, possessing no organelles, but containing hemoglobin. This hemoglobin is however homogeneously distributed towards the peripheries of red blood cells, leaving a central pallor area (30-45% of RBC

diameter) [55]. Also, the normal erythrocyte membrane appears to be smooth without any protuberances and is basically indistinct from the Hb content. There are some factors that can alter the morphology of normal RBCs such as an alteration (either increase or decrease) in Hb volume; a change in Hb structure (i.e. Hb variant, HbC, sickle cell Hb; protective enzyme deficiency leading to polymerization and precipitation); increase level of erythrocyte destruction or increase in level of demand (i.e. insufficient hemoglobinization of cells; absence or poorly functioning spleen) [56].

RBCs must maintain structural characteristics and a high amount of flexibility in order to function properly. However, a variety of abnormalities linked to certain pathologies have been described in literature and medical practice, as they were observed on peripheral blood smears. Table 2 presents, in alphabetical order, all existent such pathogenic abnormalities of RBCs (to the author's knowledge and based on available research) for simplicity of access.

Table 2. RBC pathologic variations

Name	Description	Ref.
Acanthocyte (spur cell)	Described as having long projections that are irregular (spicules). They are also characterized as having no central pallor. It is mainly associated with severe liver disease, partial or total removal of the spleen and some rare cases of abetalipoproteinemia or neuroacanthocytosis (McLeod and choreo-acanthocytosis syndrome).	[57,58]
Anisocytosis	The sizes of erythrocytes can vary within a wide range	[59-61]
Babesiosis	This appears similar to the ring forms in malaria	[62,63]
Basophilic stippling	Red blood cells present small RNA aggregates which are seen as little blue dots. In this case, fewer dots can be a characteristic of reticulocytes and more dots that of toxic bone marrow damage, thalassemia or myelodysplasia	[64,65]
Elliptocyte	The cell appears elongated and elliptical and it is associated with rare hereditary elliptocytosis	[66,67]
Echinocyte	Similar to acanthocytes but the spicules or projections are regular. Might be associated with uremia/renal disease	[58]
Heinz type body	Precipitated hemoglobin is visible after supravital staining as a perimembranous blue dot, common with hemoglobinopathies	[68,69]
Howell-Jolly type body	Commonly seen with the absence of the spleen, observed as a deeply basophilic nuclear remainder that is small and round	[70,71]
Hypochromic	Presented as cells with an increased mean corpuscular hemoglobin (MCH) and it is specific to a deficiency in iron. A RBC is considered hypochromic if the central pallor is measured at more than 50 % of the RBC diameter.	[72,73]
Macrocytosis	Red blood cells present an increased mean corpuscular volume (MCV) and it is typically associated with megaloblastic anemias. Macrocytes measure typically more than 9 μm in diameter	[74,75]

Malaria	Infections with the single-cell parasite that causes malaria can be observed as ring-forms, dots or gametocytes	[76-78]
Microcytosis	Red blood cells present a decreased mean corpuscular volume (MCV) and it is linked to iron deficiencies and thalassemia types. Microcytes measure on average less than 6 μm in diameter.	[79,80]
Pappenheimer type body	Described by the presence of little blue dots that contain iron. This is typically linked to an iron overload or to the absence of the spleen	[81,82]
Poikilocytosis	Red blood cells vary in shapes	[83]
Polychromatophilia	Young erythrocytes present a bluish tint due to high a content in RNA	[84]
Reticulocyte	Supravital staining can precipitate young RBCs with higher RNA content for identification and counting	[85,86]
Rouleaux	More serum protein, especially more fibrinogen or globulin, reduces surface charge and causes a linear aggregation of RBCs that resembles a stack of coins.	[87,88]
Sideroblast	Non-nucleated erythrocytes that have stainable iron	[89,90]
Spherocyte	RBCs appear smaller, rounder, and denser without any presence of a central pallor. This has been linked to splenic hemolysis in normal patients. In patients suffering from hereditary spherocytosis and increase in osmotic fragility can be seen	[91,92]
Schistocyte (schizocyte)	RBC shapes are presented as irregular and fragmented, and this is mainly associated with intravascular hemolysis (i.e. microangiopathic hemolytic anemias). Alternative that appears cut in half (i.e. helmet cell) or with 2, 3 pointed projections on the surface (i.e. horn cells/ keratocytes) also exist.	[93,94]
Sickle cell (drepanocyte)	Red blood cells in sickle cell pathologies are seen as curved-shaped, with the ends being pointed. This is due to aggregation of hemoglobin S	[95,96]
Stomatocyte	Erythrocytes with a central pallor that is similar to a slit, sporadically non-specific or artefactual stomatocytes. This is linked to rare hereditary stomatocytosis	[97,98]
Target cell (codocyte)	Cell with central and peripheral staining and an intervening pallor from increased RBC membrane redundancy. This can be observed in the case of some thalassemia types, liver disease, as well as hemoglobin C disease	[58, 99]
Tear drop cell (dacrocyte)	Red blood cells are pinched at one end, hence the name. These specific types of cells are linked to myelophthisic conditions and myelofibrosis	[100]
Wafer cell (leptocyte)	The hemoglobin is present at the periphery of cells which are thin and flat in appearance. Leptocytes tend to fold into bowl-shaped cells	[101]

While erythrocytes are essential for human life, due to their role in transporting oxygen, they have generally been considered inert from an immunological point of view. However, it was recently shown that they also

function as immunological sensors due to the surface expression of the nucleic acid-sensing toll-like receptor (TLR9). According to Hotz et al. erythrocytes can bind mitochondrial DNA to prevent lung injury [102]. This intrinsic immune response can act in a favorable way in the removal of damaged erythrocytes. Moreover, it also plays a part in systemic inflammation as well as the progression of anemia in pathologic states with elevated free-cell DNA.

Red blood cells are also capable of binding pathogenic organisms, the most notable of which being the *Plasmodium* species, which is associated to malaria. It is reported that different species within this genus have distinct mechanisms for adhering to RBCs [103]. For example, the most virulent malaria pathogen (*P. falciparum*) adheres to glycoporphins (A, B or C) and therefore can invade RBCs [104] to multiply inside and thus evade any repercussions from the immune system. This invasion leads to changes in erythrocytes, most notably a loss of normal shape, increased permeability to ionic species and an increased rigidity of the membrane [105].

In a recent study by Lam et al. [106] it was shown that when compared to healthy erythrocytes which have only a small percentage of TLR9 on their membrane, those from malaria patients have up to 40%. Because RBCs lack a nucleus to regulate gene expression in response to environmental factors, such changes in TLR9 protein were explained by DNA testing as being linked to changes in the membrane and shape in order to expose more of this protein. In addition, in light of the ongoing global health issue, it was discovered that the SARS-COV-2 virus can infect RBC precursors [107]. Therefore, erythrocyte production (erythropoiesis) under stress is part of the response to this virus. According to Marchi et al. [108] certain morphologies related with the infection rather than any comorbidities have also been recorded for COVID-19 individuals. Most patients in this study had spiculated cells, which could be an indicator of erythrocytes with impaired protein and lipid membrane composition.

4. RBCs and their interaction with drugs

Most drugs have potential side effects especially at higher doses or when used over extended periods of time. It is crucial to obtain the maximum therapeutic result with as few repercussions as possible. Drug activity and potency, on the other hand, are affected by a variety of parameters such as adsorption, distribution, and metabolism. Drugs are transported within the blood stream either in bound form (e.g. RBCs, plasma proteins) or in unbound form. Only the unbound form can be available for passive diffusion to sites, and typically this determines drug efficacy. Thus, the interactions of drugs with cell membranes is important to understand. However, due to the complexity of natural cell membranes, several biomimetic models have been proposed throughout the years,

for example, monolayers [109-118], bilayers [119-121] or Langmuir-Blodgett layers [122-128]. While key towards understanding interactions with drugs at a molecular level, they present certain drawbacks. Here, probably the most important one is the challenging aspect of incorporating proteins in such models, a conundrum since membrane proteins tend to have an impact on the adsorption, distribution, metabolism, and excretion of drugs. Nevertheless, such models offer valuable insights albeit their individual pros and cons.

Even so, the specific interaction of erythrocytes with intravenous injectable-type drugs (e.g. antibiotics, chemotherapy agents or insulin) is particularly important as they do interact with each other intravenously. Each drug has a different affinity to RBCs, and it is reported that there is a battle between erythrocytes and blood plasma proteins for the free drug present within the bloodstream [129]. Plasma protein binding (mostly serum albumin and α 1-acid glycoprotein) is unwanted as it can take away from drug potency [130]. Serum albumin binds mostly acidic drugs while more basic drugs are bound to lipoproteins, alpha-1 acid glycoprotein, or both [131].

In theory, the more lipophilic a medicine is, the more it binds to red blood cells; non-lipophilic substances are not dispersed to RBCs at all. Many lipophilic compounds will dissolve in RBCs lipid bilayer and then penetrate the cell. However, there are highly lipophilic substances that have been reported to have a higher protein binding affinity and thus do not necessarily bind well to RBCs. One such example is tetrahydrocannabinol that binds well (96-98%) to washed out erythrocytes (in the absence of plasma proteins) but very poorly (7-9%) to RBCs in whole blood (in the presence of plasma proteins) [132]. Substances that possess a high affinity to erythrocytes have a large blood to plasma ratio (R_b), where:

$$R_b = \frac{\text{blood concentration of drug}}{\text{plasma concentration of drug}} \quad [133]$$

The scientific literature offers substantial information on the effects of various chemicals on RBCs, with some recent research focused on changes in erythrocytes caused by drug binding. For example, Zdrengea et al. [134] investigated the effect of procaine ($5 \cdot 10^{-7}$ M, $5 \cdot 10^{-5}$ M, $5 \cdot 10^{-4}$ M) on a human erythrocyte membrane revealing changes at the micro and nanometer scales.

The aggregation of membranous particles on the cell surface and the creation of domain structures brought on by procaine are two factors that can be linked to changes in the surface morphology of the erythrocyte membrane. One other study [135] presents findings, through X-ray diffraction, on the effect of aspirin (1 mM, 1.5 mM, 2 mM, 2.5 mM, 3 mM) on the structure of RBC membranes. While the concentration of aspirin had no effect on the size of the lipid domains, there was a little increase (30 to 38 Å) in the size of the peptide domain. According to the

authors, membrane thickness is lowered while the distance between lipid tails increases, indicating membrane fluidification.

In erythrocytes, it is reported that there are 3 major components that have the capability to bind drugs, namely hemoglobin, carbonic anhydrase (intracellular enzyme that catalyzes the $\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3$ reaction and is found in RBC in high concentrations) and the actual cell membrane [129].

One study [139] found that Penicillin G, dicloxacillin, tetracycline and minocycline tend to bind not only to hemoglobin but to the globin moiety, in the absence of iron, after the dissociation of heme. Out of the four antibiotics, tetracycline stands out as it can also bind to carbonic anhydrase with zinc being required for the binding. More recent molecular docking studies [137] confirmed that certain drugs have an affinity for the specific protein moiety of hemoglobin – in this case, a lipid-lowering agent (bezafibrate) and a diuretic used for blood pressure (hydrochlorothiazide).

Another two diuretics, namely chlorthalidone and acetazolamide have been proven to bind to carbonic anhydrase. Both compete for the same binding sites and acetazolamide tends to inhibit the uptake of chlorthalidone throughout mixed treatments and being able to displace it at the binding sites [138]. Also, it was demonstrated that the accumulation process of acetazolamide within RBCs is composed of two parts, namely a non-linear one that is saturable and a linear one which is controlled by diffusion [139].

As for the RBC membrane and its interaction with drugs, for example, Bickel et al. [140] investigated the binding of an anesthetic (chlorpromazine) a tricyclic antidepressant (imipramine) and an anti-inflammatory agent (salicylic acid). It was discovered that chlorpromazine mainly binds to the membrane, although there is a possible interference as hemoglobin tends to catalyze the formation of chlorpromazine-sulphoxide. Imipramine has quite a negligible binding to erythrocyte contents, being bound mainly to its membrane. Salicylic acid on the other hand has little affinity for red blood cells, being quite partial to plasma proteins.

The examples discussed above show that drugs can behave differently, even if they are part of the same class. A better understanding of such interactions is of particular importance considering the emerging research trend of using RBCs as drug or nanoparticle delivery vehicles. Given their biocompatibility and long-term free circulation throughout the circulatory system, this is extremely possible.

These would undoubtedly extend the drug's life, which is critical for medicine that has a gradual release in blood or has a purpose of action within the intravascular area within places accessible to erythrocytes [141]. The ability to help drugs bypass metabolism mechanisms also make RBCs the ideal choice for drug delivery.

The incorporation of pharmaceuticals into red blood cells will not only guard against immune cell phagocytosis but also prevent any potential biotoxicity from the drug itself [142]. Conversely, an alternative to encapsulation is that of coupling drugs onto the erythrocyte membrane. While a concern can be raised regarding drug isolation, this is not valid when it comes to pharmaceuticals that are supposed to be active within the bloodstream [143].

5. Other considerations

While the need to study the “classic type” of injectable drugs in relation to blood components and specifically red blood cells is quite obvious, concerns can be raised in regard to other substances/devices especially pertaining to implantology. This applies even to biocompatible and bioactive types of resorbable materials. One such example would be hydroxyapatite ($\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$, HAP) a substance widely investigated for bone substitutes due to its similarity to the mineral component in human bone tissue [144-163].

Even though HAP-based implants (as a coating on metallic implants or in composites) are used at bone sites and they are technically biocompatible and bioactive, aiding in bone proliferation, there are certain things to consider. Ions (i.e. calcium and phosphate for unsubstituted HAP) can leach into the bloodstream. This can be somewhat problematic, depending on several factors such as size and type of implant or different pathologies of the patient in cause.

Intracellular calcium ions are a key factor in erythrocytes, influencing aspects including metabolic activity, redox state, and cell clearance in addition to biophysical characteristics (i.e. membrane composition, volume, and rheological characteristics) [164]. Ca^{2+} can cause K^+ efflux (Gardos effect) after it enters RBCs, causing cell degeneration (e.g., loss of intracellular fluid and volume decrease) [165,166]. However, the biomedical applications of HAP go far beyond bone implants. Due to their high biocompatibility and ability to be functionalized with different drugs and molecules, HAP nanoparticles have been the subject of a variety of targeted delivery studies.

In this case, the importance of blood compatibility studies and a deeper understanding of how these functionalized nanoparticles would interact with red blood cells and other blood components is especially important. Particulate matter within the bloodstream can hinder the homeostatic balance [167].

Even though this is not a well-investigated topic, there are some studies concentrating on hydroxyapatite and its potential impacts on red blood cells. Table 3 contains examples of some articles discovered by the authors.

Table 3. Hydroxyapatite-based materials and their effects on red blood cells

Type of HAP material	Type of study on RBCs	Effect	Ref.
HAP nanoparticles	<i>In vitro</i> 5% centrifuged RBC suspension in 0.9% NaCl solution	- aggregation and adherence to membrane - higher adsorption capacity leads to “caves” on RBC membrane - no rupture in lipid bilayer - no hemolysis	[168]
HAP nanoparticles modified with heparin		- inhibited aggregation - no hemolysis	
HAP microparticles		- no observed adhering to membrane - no hemolysis	
Hydroxyapatite - Polymethyl Methacrylate Nanocomposites (10% HAP)	<i>In vitro</i> Blood samples (healthy donors 20-28 years) were mixed with 50% (mg/dL) HAP-PMMA	- decreased RBC lifetime - change in RBC shape and size - lower hemoglobin in 40% samples	[169]
Commercial Rod-shaped HAP (width -15–30 nm, length - 40–70 nm)	<i>In vitro</i> Blood - healthy non-smoking adult volunteers Free from medication	~ 0.25% hemolysis at 1 and 10 mg/mL	[170]
Commercial Needle-shaped HAP (width - 30–60 nm, length - 200–500 nm)			
Synthetic HAP diameter: 3-11 nm length: 20-65 nm	RBC centrifuged from BALB/c mice blood 5% (v/v) RBC in saline	- lower hemolytic activity (<5%) compared to control - this may be caused by surface hydrophilic groups (carboxyl, hydroxyl functional groups)	[171]
hydroxyapatite magnetic nanoparticles doped with different ions (Gd ³⁺ / Fe ²⁺ / Fe ³⁺ / Co ²⁺)	Blood from 4 volunteers	- no hemolytic activity up to concentrations of 4 mg/mL - nanoparticles do not affect the membrane integrity or morphology of RBCs	[172]

6. Conclusions

Recent studies regarding red blood cells have revealed that their functions go beyond that of oxygen transporters, as they play a key role in redox regulation, metabolism control and interorgan communication. However, it is possible to state that they are still understudied. Because erythrocytes are essential for life, more research is needed to understand every aspect of erythrocytes, from their development in the bone marrow to their end of life and various diseases, to move forward with different clinical applications.

Acknowledgment

This work was supported by grants of the Ministry of Research, Innovation and Digitization, CNCS/CCCDI-UEFISCDI, project number 186 PCE 2021.

REFERENCES

- [1] S.J. Fathima, F. Khanum, Blood cells and leukocyte culture – a short review, *Blood Research & Transfusion Journal*, **1**(2), 555559 (2017). <http://dx.doi.org/10.19080/OABTJ.2017.01.555559>
- [2] K. Almezghwi, S. Serte, Improved classification of white blood cells with the generative adversarial network and deep convolutional neural network, *Computational Intelligence and Neuroscience*, **2020**, 6490479 (2020). <https://doi.org/10.1155/2020/6490479>
- [3] O. Leiter, T.L. Walker, Platelets: The missing link between the blood and brain?, *Progress in Neurobiology*, **183**, 101695 (2019). <https://doi.org/10.1016/j.pneurobio.2019.101695>
- [4] S. Himbert, M.C. Rheinstädter, Structural and mechanical properties of the red blood cell's cytoplasmic membrane seen through the lens of biophysics, *Frontiers in Physiology*, **13**, 953257 (2022). <https://doi.org/10.3389/fphys.2022.953257>
- [5] L.L. Munn, M.M. Dupin, Blood cell interactions and segregation in flow, *Annals of Biomedical Engineering*, **36**(4), 534–544 (2008). <https://doi.org/10.1007/s10439-007-9429-0>
- [6] V. Pretini, M.H. Koenen, L. Kaestner, M.H.A.M. Fens, R.M. Schiffelers, M. Bartels, R. Van Wijk, Red Blood cells: chasing interactions, *Frontiers in Physiology*, **10**, 945 (2019). <https://doi.org/10.3389/fphys.2019.00945>
- [7] D.Z. de Back, E.B. Kostova, M. van Kraaij, T.K. van den Berg, R. van Bruggen, Of macrophages and red blood cells; a complex love story, *Frontiers in Physiology*, **5**, 9 (2014). <https://doi.org/10.3389/fphys.2014.00009>
- [8] J.L. Vives Corrons, L. Berga Casafont, E. Feliu Fransedo, Concise review: how do red blood cells born, live, and die? *Annals of Hematology*, **100**, 2425-2433 (2021). <https://doi.org/10.1007/s00277-021-04575-z>
- [9] C. Mussolino, J. Strouboulis, Recent Approaches for manipulating globin gene expression in treating hemoglobinopathies. *Frontiers in Genome Editing*, **3**, 618111 (2021). <https://doi.org/10.3389/fgeed.2021.618111>
- [10] V. Kuhn, L. Dietrich, T.C. Stevenson Keller IV, C.M. Kramer, W. Luckstadt, C. Panknin, T. Suvorava, B.E. Isakson, M. Kelm, M.M. Cortese-Krott, Red blood cell function and dysfunction: redox regulation, nitric oxide metabolism, anemia. *Antioxidants & Redox Signaling*, **26**(13), 718-741 (2017). <https://doi.org/10.1089/ars.2016.6954>
- [11] P. Thiagarajan, C.J. Parker, J.T. Prchal, How do red blood cells die?, *Frontiers in Physiology*, **12**, 655393 (2021). <https://doi.org/10.3389/fphys.2021.655393>
- [12] H. Li, G. Lykotrafitis, Erythrocyte membrane model with explicit description of the lipid bilayer and the spectrin network, *Biophysical Journal*, **107**(3), 642-653 (2014). <http://dx.doi.org/10.1016/j.bpj.2014.06.031>
- [13] S. De Oliveira, C. Saldanha, An overview about erythrocyte membrane, *Clinical Hemorheology and Microcirculation*, **44**, 63–74 (2010). <https://doi.org/10.3233/CH-2010-1253>

- [14] A. Vahedi, P. Bigdelou, A.M. Farnoud, Quantitative analysis of red blood cell membrane phospholipids and modulation of cell-macrophage interactions using cyclodextrins, *Scientific Reports*, **10**, 15111 (2020). <https://doi.org/10.1038/s41598-020-72176-3>
- [15] S.E. Lux IV, Anatomy of the red cell membrane skeleton: unanswered questions, *Blood*, **127**(2), 187-199 (2016). <https://doi.org/10.1182/blood-2014-12-512772>
- [16] M.L. Jennings, Cell physiology and molecular mechanism of anion transport by erythrocyte band 3/AE1, *American Journal of Physiology Cell Physiology*, **321**, C1028-C1059 (2021). <https://doi.org/10.1152/ajpcell.00275.2021>
- [17] K. Hsu, Exploring the Potential Roles of Band 3 and Aquaporin-1 in Blood CO₂ Transport– Inspired by Comparative Studies of Glycophorin B-A-B Hybrid Protein GP.Mur, *Frontiers in Physiology*, **9**, 733 (2018). <https://doi.org/10.3389/fphys.2018.00733>
- [18] E.J. Hollox, S. Louzada, Genetic variation of glycophorins and infectious disease, *Immunogenetics*, (2022). <https://doi.org/10.1007/s00251-022-01280-7>
- [19] K. Giger, I. Habib, K. Ritchie, P.S. Low, Diffusion of glycophorin a in human erythrocytes, *Biochimica et Biophysica Acta*, **1858**(11), 2839-2845 (2016). <https://doi.org/10.1016%2Fj.bbamem.2016.08.012>
- [20] A. Schawalter, M.E. Reid, K. Yazdanbakhsh, Recombinant glycophorins C and D as tools for studying Gerbich blood group antigens, *Immunohematology*, **44**, 567-574 (2004).
- [21] R. Zhang, C.Y. Zhang, Q. Zhao, D.H. Li, Spectrin: Structure, function and disease, *Science China Life Sciences*, **56**(12), 1076-1085 (2013). <https://doi.org/10.1007/s11427-013-4575-0>
- [22] C. Leterrier, P.A. Pullarkat, Mechanical role of the submembrane spectrin scaffold in red blood cells and neurons, *Journal of Cell Science*, **135**, jcs259356 (2022). <https://doi.org/10.1242/jcs.259356>
- [23] E. Kozlova, A. Chernysh, V. Sergunova, E. Manchenko, V. Moroz, A. Kozlov, Conformational distortions of the red blood cell spectrin matrix nanostructure in response to temperature changes in vitro, *Scanning*, **2019**, 8218912 (2019). <https://doi.org/10.1155/2019/8218912>
- [24] D.S. Gokhin, R.B. Nowak, J.A. Khoory, A. De la Piedra, I.C. Ghiran, V.M. Fowler, Dynamic actin filaments control the mechanical behavior of the human red blood cell membrane, *Molecular Biology of the Cell*, **26**, 1699-1710 (2015). <https://doi.org/10.1091/mbc.E14-12-1583>
- [25] G.S. Gokhin, V.M. Fowler, Feisty filaments: actin dynamics in the red blood cell membrane skeleton, *Current Opinion in Hematology*, **23**(3), 206-214 (2016). <https://doi.org/10.1097/MOH.0000000000000227>
- [26] R.B. Nowak, H. Alimohamadi, K. Pestonjamas, P. Rangamani, V.M. Fowler, Nanoscale dynamics of actin filaments in the red blood cell membrane skeleton, *Molecular Biology of the Cell*, **33**(3), ar28 (2022). <https://doi.org/10.1091/mbc.e21-03-0107>
- [27] X. Xia, S. Liu, Z.H. Zhou, Structure, dynamics and assembly of the ankyrin complex on human red blood cell membrane, *Nature Structural & Molecular Biology*, **29**, 698-705 (2022). <https://doi.org/10.1038/s41594-022-00779-7>
- [28] W. Sae-Lee, C.L. McCafferty, E.J. Verbeke, P.C. Havugimana, O. Papoulas, C.D. McWhite, J.R. Houser, K. Vanuytsel, G.J. Murphy, K. Drew, A. Emili, D.W. Taylor, E.M. Marcotte, The protein organization of a red blood cell, *Cell Reports*, **40**, 111103 (2022). <https://doi.org/10.1016/j.celrep.2022.111103>
- [29] T.J. Satchwell, D.K. Shoemark, R.B. Sessions, A.M. Toye, Protein 4.2 : A complex linker, *Blood Cells, Molecules, and Diseases*, **42**, 201-210 (2009). <https://doi.org/10.1016/j.bcmed.2009.01.005>

- [30] Y. Su, Y. Ding, M. Jiang, W. Jiang, X. Hu, Z. Zhang, Associations of protein 4.2 with band 3 and ankyrin, *Molecular and Cellular Biochemistry*, **289**, 159-166 (2006). <https://doi.org/10.1007/s11010-006-9159-x>
- [31] M. Solomao, X. Zhang, Y. Yang, S. Lee, J.H. Hartwig, J.A. Chasis, N. Mohandas, X. An, Protein 4.1R-dependent multiprotein complex: New insights into the structural organization of the red blood cell membrane, *PNAS*, **105**(23), 8026-8031 (2008). www.pnas.org/cgi/doi/10.1073/pnas.0803225105
- [32] B.S. Nayak, V.Y. Beharry, S. Armoogam, M. Nancoo, K. Ramadhin, K. Ramesar, C. Ramnarine, A. Singh, A. Singh, K.U. Nwachi, S. Teelucksing, R. Mathura, L. Roberts, Determination of RBC membrane and serum lipid composition in trinidadian type II diabetics with and without nephropathy, *Vascular Health and Risk Management*, **4**(4), 893-899 (2008). <https://doi.org/10.2147/vhrm.s2656>
- [33] A. Vahedi, P. Bigdelou, A.M. Farnoud, Quantitative analysis of red blood cell membrane phospholipids and modulation of cell-macrophage interactions using cyclodextrins, *Scientific reports*, **10**, 15111 (2020). <https://doi.org/10.1038/s41598-020-72176-3>
- [34] D. Bladier, G. Perret, J. Baudelot, P. Cornillot, F. Gourbeault, Carbohydrate Content of Human Erythrocyte Membrane. Variations with ABO-Blood Group, *Hoppe-Seyler's Zeitschrift für Physiologische Chemie*, **360**, 501-506 (1979). <https://doi.org/10.1515/bchm2.1979.360.1.501>
- [35] Gh. Benga, O. Popescu, V. Borza, V.I. Pop, I. Mocsy, A. Brain, J. Wrigglesworth, Water permeability of human erythrocytes. Identification of membrane proteins involved in water transport, *European Journal of Cell Biology*, **41**(2), 252-262 (1986).
- [36] Gh. Benga, O. Popescu, V.I. Pop, R.P. Holmes, p-(Chloromercuri)benzenesulfonate binding by membrane proteins and the inhibition of water transport in human erythrocytes, *Biochemistry*, **25**(7), 1535-1538 (1986). <https://doi.org/10.1021/bi00355a011>
- [37] Gh. Benga, Water exchange through the erythrocyte membrane, *International Review of Cytology*, **114**, 273-316 (1989). [https://doi.org/10.1016/S0074-7696\(08\)60864-5](https://doi.org/10.1016/S0074-7696(08)60864-5)
- [38] Gh. Benga, Water transport in human red blood cells, *Progress in Biophysics and Molecular Biology*, **51**, 193-245 (1988).
- [39] Gh. Benga, The first discovered water channel protein, later called aquaporin 1: Molecular characteristics, functions and medical implications, *Molecular Aspects of Medicine*, **33**, 518-534 (2012). <http://dx.doi.org/10.1016/j.mam.2012.06.001>
- [40] B.M. Denker, B.L. Smith, F.P. Kuhajda, P. Agre, Identification, purification and partial characterization of a novel Mr 28,000 integral membrane protein from erythrocytes and renal tubules, *Journal of Biological Chemistry*, **263**, 15634-15642 (1988).
- [41] M.L. Zeidel, S.V. Ambudkar, B.L. Smith, P. Agre, Reconstitution of functional water channel in liposomes containing purified red cell CHIP28 protein, *Biochemistry*, **31**, 7436-7440 (1992).
- [42] G.M. Preston, T.P. Carroll, W.B. Guggino, P. Agre, Appearance of water channels in *Xenopus* oocytes expressing red cell CHIP 28 protein, *Science*, **256**, 385-387 (1992).
- [43] C. Sut, S. Tariket, M.L. Chou, O. Garraud, S. Laradi, H. Hamzeh-Cognasse, J. Seghatchian, T. Burnouf, F. Cognasse, Duration of red blood cell storage and inflammatory marker generation, *Blood Transfusion*, **15**(2), 145-152 (2017). <https://doi.org/10.2450%2F2017.0343-16>
- [44] M.S.Y. Ng, M. David, R.A. Middelburg, A.S.Y. Ng, J.Y. Suen, J.P. Tung, J.F. Fraser, Transfusion of packed red blood cells at the end of shelf life is associated with increased risk of mortality – a pooled patient data analysis of 16 observational trials, *Hematologica*, **103**(9), 1542-1548 (2018). <https://doi.org/10.3324%2Fhaematol.2018.191932>

- [45] J. Lecak, K. Scott, C. Young, J. Hannon, J.P. Acker, Evaluation of red blood cells stored at -80 °C in excess of 10 years, *Blood Components*, **44**, 1306-1313 (2004). <https://doi.org/10.1111/j.1537-2995.2004.03271.x>
- [46] A. Chang, Y. Kim, R. Hoehn, P. Jernigan, T. Pritts, Cryopreserved packed red blood cells in surgical patients: past, present, and future, *Blood Transfusion*, **15**(4), 341-347 (2017). <https://doi.org/10.2450%2F2016.0083-16>
- [47] D.A. Hampton, C. Wiles, L.J. Fabricant, L. Kiraly, J. Differding, S. Underwood, D. Le, J. Watters, M.A. Schreiber, Cryopreserved red blood cells are superior to standard liquid red blood cells, *Journal of Trauma and Acute Cell Surgery*, **77**(1), 2027 (2014). <https://doi.org/10.1097/TA.0000000000000268>
- [48] R.H. Bremmer, K.G. de Bruin, J.C. van Gemert, T.G. van Leeuwen, M.C.G. Aalders, Forensic quest for age determination of bloodstains, *Forensic Science International*, **216**, 1-11 (2012). <https://doi.org/10.1016/j.forsciint.2011.07.027>
- [49] M. Tsuruga, A. Matsuoka, A. Hachimori, Y. Sugawara, K. Shikama, The molecular mechanism of autoxidation for human oxyhemoglobin. Tilting of the distal histidine causes nonequivalent oxidation in the beta chain, *Journal of Biological Chemistry*, **273**(15), 8607-8615 (1998). <https://doi.org/10.1074/jbc.273.15.8607>
- [50] H. Lin, Y. Zhang, Q. Wang, B. Li, P. Huang, Z. Wang, Estimation of the age of human bloodstains under the simulated indoor and outdoor crime scene conditions by ATR-FTIR spectroscopy, *Scientific reports*, **7**, 13254 (2017). <https://doi.org/10.1038/s41598-017-13725-1>
- [51] G. Edelman, T.G. van Leeuwen, M.C.G. Aalders, Hyperspectral imaging for the age estimation of blood stains at the crime scene, *Forensic Science International*, **223**, 72-77 (2012). <http://dx.doi.org/10.1016/j.forsciint.2012.08.003>
- [52] S. Strasser, A. Zink, G. Kada, P. Hinterdorfer, O. Peschel, W.M. Heckl, A.G. Nerlich, S. Thalhammer, Age determination of blood spots in forensic medicine by force spectroscopy, *Forensic Science International*, **170**, 8-14 (2007). <http://dx.doi.org/10.1016/j.forsciint.2006.08.023>
- [53] D.R. Cavalcanti, L.P. Silva, Application of atomic force microscopy in the analysis of time since deposition (TDS) of red blood cells in bloodstains: a forensic analysis, *Forensic Science International*, **301**, 254-262 (2019). <https://doi.org/10.1016/j.forsciint.2019.05.048>
- [54] T. Smijs, A. Hosseinzoi, F. Galli, Forensic application of atomic force microscopy for age determination of bloodstains, *Journal of Forensic Investigation*, **5**(1), 6 (2017).
- [55] E.C. Lynch, Peripheral Blood Smear. In: Walker HK, Hall WD, Hurst JW, editors. *Clinical Methods: The History, Physical, and Laboratory Examinations*. 3rd edition. Boston: Butterworths; 1990. Chapter 155. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK263>
- [56] L. Tyrell, G. Rose, A. Shukri, S.B. Kahwash, Morphologic changes in red blood cells: An illustrated review of clinically important light microscopic findings, *The Malaysian Journal of Pathology*, **43**(2), 219-239 (2021).
- [57] J.A. McBride, H.S. Jacob, Abnormal kinetics of red cell membrane cholesterol in acanthocytes: studies in genetic and experimental abetalipoproteinaemia and in spur cell anemia, *British Journal of Haematology*, **18**, 383-397 (1970). <https://doi.org/10.1111/j.1365-2141.1970.tb01452.x>
- [58] R. Sharma, C.J. Holman, K.E. Brown, A thorny matter: Spur cell anemia, *Annals of Hepatology*, **28**, 100771 (2023). <http://creativecommons.org/licenses/by-nc-nd/4.0/>
- [59] G. Nicolas, M. Bennoun, A. Porteu, S. Mativet, C. Beaumont, B. Grandchamp, M. Sirito, M. Sawadogo, A. Kahn, S. Vaulont, PNAS, **99**(7), 4596-4601 (2002). <https://doi.org/10.1073/pnas.072632499>

- [60] G.S. Sultana, S.A. Haque, T. Sultana, Q. Rahman, A.N.N. Ahmed, Role of red cell distribution width (RDW) in the detection of iron deficiency anaemia in pregnancy within the first 20 weeks of gestation. *Bangladesh Medical Research Council Bulletin*, **37**(3), 102–105 (2011). <https://doi.org/10.3329/bmrcb.v37i3.9122>
- [61] D. Kust, M. Lucijanac, K. Urch, I. Samija, I. Celap, I. Kruljac, M. Prpic, I. Lucijanac, N. Matesa, A. Bolanca, Clinical and prognostic significance of anisocytosis measured as a red cell distribution width in patients with colorectal cancer, *QJM: An International Journal of Medicine*, **110**(6), 361-367 (2017). <https://doi.org/10.1093/qjmed/hcw223>
- [62] M.J. Homer, I. Aguilar-Delfin, S.R. Telford III, P.J. Krause, D.H. Persing, Babesiosis, *American Society for Microbiology, Clinical Microbiology Reviews*, **13**(3), 451-469 (2000). <https://doi.org/10.1128/CMR.13.3.451>
- [63] P.J. Krause, Human Babesiosis, *International Journal of Parasitology*, **49**(2), 165-174 (2019). <https://doi.org/10.1016/j.ijpara.2018.11.007>
- [64] J.-L. Vives Corrons, B.J. Bain, Haemoglobin Bristol-Alesha in a child with non-spherocytic severe haemolytic anaemia and marked anisochromic poikilocytosis with basophilic stippling and amorphous intracellular content, *Blood Cells, Molecules and Diseases*, **94**, 102652 (2022). <https://doi.org/10.1016/j.bcmed.2022.102652>
- [65] B. Santosa, H.W. Subagio, L. Suromo, H.R. Sunoko, Zinc supplementation decreases basophilic stippling in rats exposed to lead, *Universa Medicina*, **33**(1), 11-18 (2014).
- [66] A. Vaya, M. Suescun, A. Pardo, O. Fuster, Erythrocyte deformability and hereditary elliptocytosis, *Clinical Hemorheology and Microcirculation* **58**, 471–473 (2014). <https://doi.org/10.3233/ch-141889>
- [67] D. Sidam, P. Gogoi, W.T. Murry, Hereditary Elliptocytosis- A Case Report, *World Wide Journal of Multidisciplinary Research and Development*, **3**(8), 188-190 (2017).
- [68] N. Bexfield, J. Archer, M. Herrtage, Heinz body haemolytic anaemia in a dog secondary to ingestion of a zinc toy: A case report, *The Veterinary Journal*, **174**(2), 414-417 (2007). <https://doi.org/10.1016/j.tvjl.2006.08.015>
- [69] C. Baetge, L.C. Smith, C.P. Azevedo, Clinical heinz body anemia in a cat after repeat propofol administration case report, *Frontiers in Veterinary Science*, **7**, 591556 (2020). <https://doi.org/10.3389/fvets.2020.591556>
- [70] D.A. Sears, M.M. Udden, Howell-Jolly bodies: a brief historical review, *The American journal of Medical Sciences*, **343**(5), 407-409 (2012). <https://doi.org/10.1097/MAJ.0b013e31823020d1>
- [71] V.L. Harrod, T.A. Howard, S.A. Zimmerman, S.D. Dertinger, R.E. Ware, Quantitative analysis of Howell-Jolly bodies in children with sickle cell disease, *Experimental Hematology*, **35**(2), 179-183 (2007). <https://doi.org/10.1016/j.exphem.2006.09.013>
- [72] W.C. Winkelmayer, M. Lorenz, R. Kramar, W.H. Horl, Sunder-Plassmann, Percentage of hypochromic red blood cells is an independent risk factor for mortality in kidney transplant recipients, *American Journal of Transplantation*, **4**(12), 2075-2081 (2004). <https://doi.org/10.1046/j.1600-6143.2004.00604.x>
- [73] M. Rehu, S. Ahonen, K. Punnonen, The diagnostic accuracy of the percentage of hypochromic red blood cells (%HYPOm) and cellular hemoglobin in reticulocytes (CHr) in differentiating iron deficiency anemia and anemia of chronic diseases, *Clinica Chimica Acta*, **412**(19-20), 1809-1813 (2011). <https://doi.org/10.1016/j.cca.2011.06.004>
- [74] F. Aslinia, J.J. Mazza, S.H. Yale, Megaloblastic anemia and other causes of Macrocytosis, *Clinical Medicine & Research*, **4**(3), 236-241 (2006). <https://doi.org/10.3121/cmr.4.3.236>
- [75] G.J. Hildick-Smith, J.D. Cooney, C. Garone, L.S. Kremer, T.B. Haack, J.N. Thon, N. Miyata, D.S. Lieber, S.E. Calvo, H.O. Akman, Y.Y. Yien, N.C. Huston, D.S. Branco, D.I. Shah, M.L. Freedman, C.M. Koehler, J.E. Italiano Jr., A. Merckenschlager, S. Beblo, T.M. Strom, T. Meitinger, P. Freisinger, M.A. Donati, H. Prokisch, V.K. Mootha, S. DiMauro, B.H.

- Paw, Macrocytic Anemia and mitochondriopathy resulting from a defect in Sideroflexin 4, *The American Journal of Human Genetics*, **93**, 906–914 (2013). <http://dx.doi.org/10.1016/j.ajhg.2013.09.011>
- [76] N. Mohandas, X. An, Malaria and human red blood cells, *Medical Microbiology and Immunology*, **201**, 593–598 (2012). <https://doi.org/10.1007/s00430-012-0272-z>
- [77] T.N. Williams, Human red blood cell polymorphisms and malaria, *Current Opinion in Microbiology*, **9**(4), 388–394 (2006). <https://doi.org/10.1016/j.mib.2006.06.009>
- [78] B.M. Cooke, N. Mahondas, R.L. Coppel, Malaria and the red blood cell membrane, *Seminars in Hematology*, **41**(2), 173–188 (2004). <https://doi.org/10.1053/j.seminhematol.2004.01.004>
- [79] E. Urrechaga, Red blood cell microcytosis and hypochromia in the differential diagnosis of iron deficiency and β -thalassaemia trait, *International Journal of Laboratory Hematology*, **31**(5), 528–534 (2009). <https://doi.org/10.1111/j.1751-553X.2008.01073.x>
- [80] M. Schoorl, M. Schoorl, J. van Pelt, P.C.M. Bartels, Application of innovative hemocytometric parameters and algorithms for improvement of microcytic anemia discrimination, *Hematology Reports*, **7**(2), 5843 (2015). <https://doi.org/10.4081/hr.2015.5843>
- [81] D.A. Sears, M.M. Udden, Pappenheimer bodies: A brief historical review, *American Journal of Hematology*, **75**(4), 249–250 (2004). <https://doi.org/10.1002/ajh.20008>
- [82] T. Inaba, A. Ohama, Prominent increase of Pappenheimer body-containing erythrocytes in a patient with hypoplastic spleen (IJHM-D-18-00279R2), *International Journal of Hematology*, **108**, 351–352 (2018). <https://doi.org/10.1007/s12185-018-2512-5>
- [83] M.M. Christopher, M.G. Hawkins, A.G. Burton, Poikilocytosis in rabbits: prevalence, type, and association with disease, *PlosOne*, **9**(11), e112455 (2014). <https://doi.org/10.1371/journal.pone.0112455>
- [84] L.A. Mathias, T.C. Fisher, L. Zeng, H.J. Meiselman, K.I. Weinberg, A.L. Hiti, Ineffective erythropoiesis in β -thalassemia major is due to apoptosis at the polychromatophilic normoblast stage, *Experimental Hematology*, **28**(12), 1343–1353 (2000). [https://doi.org/10.1016/S0301-472X\(00\)00555-5](https://doi.org/10.1016/S0301-472X(00)00555-5)
- [85] L. Blanc, A. De Gassart, C. Geminard, P. Bette-Bobillo, M. Vidal, Exosome release by reticulocytes—An integral part of the red blood cell differentiation system, *Blood Cells, Molecules, and Diseases*, **35**(1), 21–26 (2005). <https://doi.org/10.1016/j.bcmd.2005.04.008>
- [86] P. Oustamanolakis, I.E. Koutroubakis, I. Messaritakis, G. Kefalogiannis, M. Niniraki, E.A. Kouroumalis, Measurement of reticulocyte and red blood cell indices in the evaluation of anemia in inflammatory bowel disease, *Journal of Crohn's and Colitis*, **5**(4), 295–300 (2011). <https://doi.org/10.1016/j.crohns.2011.02.002>
- [87] C. Wagner, P. Steffen, S. Svetina, Aggregation of red blood cells: From rouleaux to clot formation, *Comptes Rendus Physique*, **14**(6), 459–469 (2013). <https://doi.org/10.1016/j.crhy.2013.04.004>
- [88] T. Wang, T.-W. Pan, Z.W. Xing, R. Glowinski, Numerical simulation of rheology of red blood cell rouleaux in microchannels, *Physical Review E*, **79**, 041916 (2009). <https://doi.org/10.1103/PhysRevE.79.041916>
- [89] C. Jouzier, A. Cherait, P. Cony-Makhoul, J.-F. Hamel, M. Veloso, S. Thepot, T. Cluzeau, A. Stamatoullas, A. Garnier, A. Guerci-Bresler, S. Dimicoli-Salazar, G.M. Pica, S. Cheze, C. Santana, F. Chermat, P. Fenaux, S. Park, Red blood cell transfusion burden in myelodysplastic syndromes (MDS) with ring Sideroblasts (RS): A retrospective multicenter study by the Groupe Francophone des Myélodysplasies (GFM), *Transfusion*, **62**(5), 961–973 (2022). <https://doi.org/10.1111/trf.16884>

- [90] L. Malcovanti, L. Cazzola, Refractory anemia with ring sideroblasts, *Best Practice & Research Clinical Haematology*, **26**(4), 377-385 (2013). <https://doi.org/10.1016/j.beha.2013.09.005>
- [91] E.B. Roll, T. Christensen, P.A. Gederaas, Effects of bilirubin and phototherapy on osmotic fragility and haematoporphyrin-induced photohaemolysis of normal erythrocytes and Spherocytes, *Acta Paediatrica*, **94**, 1443-1447 (2005). <https://doi.org/10.1080/08035250510032637>
- [92] H.M. Yaish, R.D. Christensen, A. Agarwal, A neonate with Coombs-negative hemolytic jaundice with spherocytes but normal erythrocyte indices: a rare case of autosomal-recessive hereditary spherocytosis due to alpha-spectrin deficiency, *Journal of Perinatology*, **33**, 404-406 (2013). <https://doi.org/10.1038/jp.2012.67>
- [93] J.F. Lesesve, S. Salignac, F. Alla, M. Defente, M. Benbih, P. Bordigoni, T. Lecompte, Comparative evaluation of schistocyte counting by an automated method and by microscopic determination. *American Journal of Clinical Pathology*, **121**(5), 739-745 (2004). <https://doi.org/10.1309/my70-7798-9kwd-yp88>
- [94] J.-F. Lesesve, V. Asnafi, F. Braun, G. Zini, Fragmented red blood cells automated measurement is a useful parameter to exclude schistocytes on the blood film, *International Journal of Laboratory Hematology*, **34**, 566-576 (2012). <https://doi.org/10.1111/j.1751-553X.2012.01434.x>
- [95] A.P. Fellows, M.T.L. Casford, P.B. Davies, J.S. Gibson, J.N. Brewin, D.C. Rees, Nanoscale adhesion profiling and membrane characterisation in sickle cell disease using hybrid atomic force microscopy-IR spectroscopy, *Colloids and Surfaces B: Biointerfaces*, **197**, 111383 (2021). <https://doi.org/10.1016/j.colsurfb.2020.111383>
- [96] K. Wang, Z. Li, O. Egini, R. Wadgaonkar, X.-C. Jiang, Y. Chen, Atomic force microscopy reveals involvement of the cell envelope in biomechanical properties of sickle erythrocytes, *BMC Biology*, **21**, 31 (2023). <https://doi.org/10.1186/s12915-023-01523-3>
- [97] D. Darghouth, B. Koehl, J.F. Heilier, G. Madalinski, P. Bovee, G. Bosman, J. Delaunay, C. Junot, P.-H. Romeo, Alterations of red blood cell metabolome in overhydrated hereditary stomatocytosis, *Hematologica*, **96**(12), 1861-1865 (2011). <https://doi.org/10.3324%2Fhaematol.2011.045179>
- [98] J.F. Flatt, L.J. Bruce, The molecular basis for altered cation permeability in hereditary stomatocytic human red blood cells, *Frontiers in Physiology*, **9**, 367 (2018). <https://doi.org/10.3389/fphys.2018.00367>
- [99] P. Chaichompoo, A. Qillah, P. Sirankapracha, J. Kaewchuchuen, P. Rimthong, K. Paiboonsukwong, Abnormal red blood cell morphological changes in thalassaemia associated with iron overload and oxidative stress, *Journal of Clinical Pathology*, **72**, 520-524 (2019). <http://dx.doi.org/10.1136/jclinpath-2019-205775>
- [100] S. Rojas-Maya, L.A. Sarias-Cueto, I. Perez-Dias, H.K. Osorio-Landa, B. Garcia-Martinez, R. Fagundo-Sierra, R. Rivera-Moscoso, R. Carrillo-Moscoso, E. Carrillo-Maravilla, S.C. Laguna-Barcenas, Prevalence of dacrocytosis in patients with chronic diseases: splenomegaly is not mandatory for teardrop cells genesis, *Journal of bone Research*, **4**(2), 167 (2016). <https://doi.org/10.4172/2572-4916.1000167>
- [101] R.K. Marganwar, V.T. Dhurvey, Human red blood cell abnormalities in fluoride endemic area of Warora Tehsil, Chandrapur District, Maharashtra, India, *International Journal of Zoological Investigations*, **7**(2), 456-461 (2021). <https://doi.org/10.33745/ijzi.2021.v07i02.020>
- [102] M.J. Hotz, D. Qing, M.G.S. Shashaty, P. Zhang, H. Faust, N. Sondheimer, S. Rivella, G.S. Worthen, N.S. Mangalmurti, Red Blood Cells Homeostatically Bind Mitochondrial DNA through TLR9 to Maintain Quiescence and to Prevent Lung Injury. *American Journal of*

- Respiratory and Critical Care Medicine, **197**, 470–480 (2018).
<https://doi.org/10.1164/rccm.201706-1161OC>
- [103] H.L. Anderson, I.E. Bridsky, N.S. Mangalmurti, The evolving erythrocyte: red blood cells as modulators of innate immunity, *The Journal of Immunology*, **201**(5), 1343-1351 (2018).
<https://doi.org/10.4049/jimmunol.1800565>
- [104] A.F. Cowman, D. Berry, J. Baum, The cellular and molecular basis for malaria parasite invasion of the human red blood cell, *Journal of Cell Biology*, **198**, 961–971 (2012).
- [105] A.K. Dasanna, S. Hillringhaus, G. Gompfer, D.A. Fedosov, Effect of malaria parasite shape on its alignment at erythrocyte membrane, *ELife*, **10**, e68818 (2021).
<https://doi.org/10.7554/eLife.68818>
- [106] L.K.M. Lam, S. Murphy, D. Kokkinaki, A. Venosa, S. Sherrill-Mix, C. Casu, S. Rivella, A. Weiner, J. Park, S. Shin, A. Vaughan, B.H. Hahn, A.R.O. John, N.J. Meyer, C.A. Hunter, G.S. Worthen, N.S. Mangalmurti, DNA binding to TLR9 expressed by red blood cells promotes innate immune activation and anemia, *Science Translational Medicine*, **13**(616), eabj1008 (2021). <https://doi.org/10.1126/scitranslmed.abj1008>
- [107] S. Shahbaz, L. Xu, M. Osman, W. Shigl, J. Shields, M. Joyce, D.L. Tyrrell, O. Oyegbami, S. Elahi, Erythroid Precursors and Progenitors Suppress Adaptive Immunity and Get Invaded by SARS-CoV-2, *Stem Cell Reports*, **16**(5), 1165–1181 (2021).
<https://doi.org/10.1016/j.stemcr.2021.04.00>
- [108] G. Marchi, L. Bozzini, F. Dima, F. Busti, A. Castagna, C. Stranieri, A.M. Frata Pasini, S. Friso, G. Lippi, D. Girelli, A. Vianello, Red blood cell morphologic abnormalities in patients hospitalized for COVID-19, *Frontiers in Physiology*, **13**, 932013 (2022).
<https://doi.org/10.3389/fphys.2022.932013>
- [109] A. Mocanu, P.J. Quinn, C. Nicula, S. Riga, Gh. Tomoaia, C.-A. Bardas, M. Tomoaia-Cotisel, Interaction of local anesthetic procaine with phospholipid model membrane, *Revue Roumaine de Chimie*, **66**(10-11), 855-869 (2021).
<https://doi.org/10.33224/rch.2021.66.10-11.09>
- [110] M. Tomoaia-Cotisel, J. Zsako, A. Mocanu, M. Lupea, E. Chifu, Insoluble mixed monolayers. III. The ionization characteristics of some fatty acids at the air/water interface, *Journal of Colloid and Interface Science*, **117**(2), 464-476 (1987).
- [111] M. Tomoaia-Cotisel, J. Zsako, E. Chifu, Dipalmitoyl lecithin and egg lecithin monolayers at an air/water interface, *Annali di Chimica (Rome)*, **71**(3-4), 189-200 (1981).
- [112] J. Zsako, M. Tomoaia-Cotisel, A. Mocanu, E. Chifu, Insoluble mixed monolayers. II. Protolytic equilibria and the influence of the pH on the collapse pressure, *Journal of Colloid and Interface Science*, **110**(2), 317-334 (1986).
- [113] M. Tomoaia-Cotisel, A. Sen, P. J. Quinn, Surface active properties of 1,2-distearoylgalactosylglycerols, *Journal of Colloid and Interface Science*, **94**, 390-398 (1983).
- [114] M. Tomoaia-Cotisel, J. Zsako, E. Chifu, P. J. Quinn, Influence of electrolytes on the monolayers properties of saturated galactolipids at the air/water interface, *Chemistry and Physics of Lipids*, **34**(1), 55-64 (1983).
- [115] P.T. Frangopol, D. A. Cadenhead, Gh. Tomoaia, A. Mocanu, M. Tomoaia-Cotisel, The effect of procaine on lipid domains investigated by contact mode atomic force microscopy, *Revue Roumaine de Chimie*, **60**(2-3), 265-273 (2015).
- [116] P.T. Frangopol, D. A. Cadenhead, M. Tomoaia-Cotisel, A. Mocanu, Procaine effects on surface topography of spread dipalmitoylphosphatidylcholine monolayers, *Studia UBB Chemia*, **54**(1), 23-35 (2009).
- [117] Gh. Tomoaia, A. Tomoaia-Cotisel, M. Tomoaia-Cotisel, A. Mocanu, Kinetic study of adsorption of some biocompounds at the oil/water interface, *Central European Journal of Chemistry*, **3**(2), 347-360 (2005).

- [118] M. Tomoaia-Cotisel, R.-D. Pasca, O. Horovitz, A. Mocanu, Surface potentials of cholesterol and dimyristoyl phosphatidylcholine monolayers at the air/water interface, *Revue Roumaine de Chimie*, **56**(10-11), 1047-1053 (2011).
- [119] M. Tomoaia-Cotisel, I.W. Levin, Thermodynamic study of the effects of ursodeoxycholic acid and ursodeoxycholate on aqueous dipalmitoyl phosphatidyl choline bilayer dispersions, *The Journal of Physical Chemistry B*, **101**(42), 8477-8485 (1997).
- [120] J. Zsako, M. Tomoaia-Cotisel, E. Chifu, Insoluble mixed monolayers. I. Phase equilibria at the collapse of binary monolayers at gas/liquid interfaces, *Journal of Colloid and Interface Science*, **102**(1), 186-205 (1984).
- [121] J. Zsako, M. Tomoaia-Cotisel, E. Chifu, Insoluble mixed monolayers. V. Molecular associations in binary films: The regular association approach, *Journal of Colloid and Interface Science*, **146**(2), 353-362 (1991).
- [122] M. Tomoaia-Cotisel, V. D. Pop, Gh. Tomoaia, A. Mocanu, Cs. Racz, C.R. Ispas, O. Pascu, O.C. Borostean, Atomic force microscopy studies of Langmuir-Blodgett films. 4. The influence of aluminum substrate on dipalmitoyl phosphatidylcholine nanolayers, *Studia UBB Chemia*, **50**(1), 23-37 (2005).
- [123] M. Tomoaia-Cotisel, Gh. Tomoaia, V.-D. Pop, A. Mocanu, O. Cozar, N. Apetroaei, Gh. Popa, Atomic force microscopy studies of Langmuir-Blodgett films. 3. Phase behaviour of dipalmitoyl phosphatidyl choline monolayers, *Revue Roumaine de Chimie*, **50**(6), 471-478 (2005).
- [124] M. Tomoaia-Cotisel, L.C. Stewart, M. Kates, J. Zsako, E. Chifu, A. Mocanu, P.T. Frangopol, L.J. Noe, P.J. Quinn, Acid dissociation constants of diphytanyl glycerol phosphorylglycerol-methylphosphate, and diphytanyl glycerol phosphoryl glycerophosphate and its deoxy analog, *Chemistry and Physics of Lipids*, **100**, 41-54 (1999).
- [125] M. Tomoaia-Cotisel, Gh. Tomoaia, V.-D. Pop, A. Mocanu, O. Cozar, N. Apetroaei, Gh. Popa, Atomic force microscopy studies of Langmuir-Blodgett films. The effect of some drugs on dipalmitoyl phosphatidylcholine, *Studia UBB Physica*, **49**(3), 141-152 (2004).
- [126] I. Cojocar, A. Tomoaia-Cotisel, A. Mocanu, T. Yupsanis, M. Tomoaia-Cotisel, The effects of globular protein from aleurone cells of barley on stearic acid monolayers, *Revista de Chimie*, **68**(7), 1470-1475 (2017).
- [127] L.J. Noe, M. Tomoaia-Cotisel, M. Casstevens, P. N. Prasad, Characterization of Langmuir-Blodgett films of 3,4-didecyloxy-2,5-di(4-nitrophenylazomethine) thiophene in a stearic acid matrix, *Thin Solid Films*, **208**, 274-279 (1992).
- [128] M. Tomoaia-Cotisel, A. Tomoaia-Cotisel, T. Yupsanis, Gh. Tomoaia, I. Balea, A. Mocanu, Cs. Racz, Coating layers of major storage protein from aleurone cells of barley studied by atomic force microscopy, *Revue Roumaine de Chimie*, **51**(12), 1181-1185 (2006).
- [129] M. Ehrnebo, "Drug binding to blood cells." *Drug-protein binding*, Praeger Publishing, New York (1986): 128-137.
- [130] G.L. Trainor, The importance of plasma protein binding in drug discovery, *Expert Opinion on Drug Discovery*, **2**(1), 51-74 (2001). <https://doi.org/10.1517/17460441.2.1.51>
- [131] H. Ahmed, F. Bergmann, M. Zeitlinger, Protein binding in translational antimicrobial development-focus on interspecies differences, *Antibiotics*, **11**, 923 (2022). <https://doi.org/10.3390/antibiotics11070923>
- [132] M. Widman, S. Agurell, M. Ehrnebo, G. Jones, Binding of (+)- and (-)- Δ^1 -tetrahydrocannabinols and (-)-7-hydroxy- Δ^1 -tetrahydrocannabinol to blood cells and plasma proteins in man, *Journal of Pharmacy and Pharmacology*, **26**(11), 914-916 (1974). <https://doi.org/10.1111/j.2042-7158.1974.tb09207.x>

- [133] H. Mamada, K. Iwamoto, Y. Nomura, Y. Uesawa, Predicting blood-to-plasma concentration ratios of drugs from chemical structures and volumes of distribution in humans, *Molecular Diversity*, **25**, 1261-1270 (2021). <https://doi.org/10.1007/s11030-021-10186-7>
- [134] U.V. Zdrengea, Gh. Tomoaia, D.-V. Pop-Toader, A. Mocanu, O. Horovitz, M. Tomoaia-Cotisel, Procaine effect on human erythrocyte membrane explored by atomic force microscopy, *Combinatorial Chemistry & High Throughput Screening*, **14**(4), 237-247 (2011). <https://doi.org/10.2174/138620711795222491>
- [135] S. Himbert, R.J. Alsop, M. Rose, L. Hertz, A. Dhaliwal, J.M. Moran-Mirabal, C.P. Verschoor, D.M.E. Bowdish, L. Kaestner, C. Wagner, M.C. Rheinstädter, The molecular structure of human red blood cell membranes from highly oriented, solid supported multi-lamellar membranes, *Scientific Reports*, **7**, 39661 (2017). <https://doi.org/10.1038/srep39661>
- [136] M.L. Kornguth, C.M. Kunin, Binding of antibiotics to the human intracellular erythrocyte proteins hemoglobin and carbonic anhydrase, *The Journal of Infectious Diseases*, **133**(2), 185-193 (1976). <https://doi.org/10.1093/infdis/133.2.185>
- [137] B.L. Melo de Assis, A. Dillmann Groto, V.H. Sales da Mota, G.C. Peiter, K.N. Teixeira, Drug-protein in vitro and in silico interaction analysis with hemoglobin as a study model, *Medicina (Ribeirão Preto)*, **54**(1), e172174 (2021). <https://doi.org/10.11606/issn.2176-7262.rmrp.2021.172174>
- [138] B. Beerman, K. Hellstrom, B. Lindstrom, A. Rosen, Binding-site interaction of chlorthalidone and acetazolamide, two drugs transported by red blood cells, *Clinical Pharmacology & Therapeutics*, **17**(4), 424-432 (1975). <https://doi.org/10.1002/cpt1975174424>
- [139] S.M. Wallace, S. Riegelman, Uptake of acetazolamide by human erythrocytes in vitro, *Journal of Pharmaceutical Sciences*, **66**(5), 729-731 (1977). <https://doi.org/10.1002/jps.2600660532>
- [140] M.H. Bickel, Binding of chlorpromazine and imipramine to red cells, albumin, lipoproteins and other blood components, *Journal of Pharmacy and Pharmacology*, **27**(10), 733-738 (1975). <https://doi.org/10.1111/j.2042-7158.1975.tb09392.x>
- [141] C.H. Villa, A.C. Anselmo, S. Mitragotri, V. Muzykantov, Red blood cells: Supercarriers for drugs, biologicals, and nanoparticles and inspiration for advanced delivery systems, *Advanced Drug Delivery Reviews*, **106A**(15), 88-103 (2016). <https://doi.org/10.1016/j.addr.2016.02.007>
- [142] S. Wang, K. Han, S. Ma, X. Qi, L. Guo, X. Li X, Blood cells as supercarrier systems for advanced drug delivery, *Medicine in Drug Discovery*, **13**, 100119 (2022). <https://doi.org/10.1016/j.medidd.2021.100119>
- [143] V.R. Muzykantov, Drug delivery by red blood cells: vascular carriers designed by Mother Nature, *Expert Opinion on Drug Delivery*, **7**(4), 403-427 (2010). <https://doi.org/10.1517%2F17425241003610633>
- [144] Gh. Tomoaia, M. Tomoaia-Cotisel, L.-B. Pop, A. Pop, O. Horovitz, A. Mocanu, N. Jumate, L.-D. Bobos, Synthesis and characterization of some composites based on nanostructured phosphates, collagen and chitosan, *Revue Roumaine de Chimie*, **56**(10-11), 1039-1046 (2011).
- [145] Gh. Tomoaia, O. Soritau, M. Tomoaia-Cotisel, L.-B. Pop, A. Pop, A. Mocanu, O. Horovitz, L.-D. Bobos, Scaffolds made of nanostructured phosphates, collagen and chitosan for cell culture, *Powder Technology*, **238**, 99-107 (2013). <https://doi.org/10.1016/j.powtec.2012.05.023>
- [146] C. Prejmorean, M. Tomoaia-Cotisel, E. Vasile, G. Furtos, L. Pop, M. Moldovan, C. Sarosi, I. Petean, Characterisation of surface organisation and morphology of some new

- experimental dental resin-based composites, *International Journal of Nano and Biomaterials*, **3**(4), 344-359 (2011). <https://doi.org/10.1504/IJNBM.2011.045886>
- [147] Gh. Tomoaia, A. Mocanu, I. Vida-Simiti, N. Jumate, L.-D. Bobos, O. Soritau, M. Tomoaia-Cotisel, Silicon effect on the composition and structure of the composition and structure of nanocalcium phosphates in vitro biocompatibility to human osteoblasts, *Materials Science & Engineering C: Materials for Biological Applications*, **37**, 37-47 (2014). <https://doi.org/10.1016/j.msec.2013.12.027>
- [148] E. Forizs, F. Goga, A. Avram, A. Mocanu, I. Petean, O. Horovitz, M. Tomoaia-Cotisel, Thermal analysis of pure and multisubstituted hydroxyapatite pastes, *Studia UBB Chemia*, **62**(4), 173-180 (2017). <https://doi.org/10.24193/subbchem.2017.4.14>
- [149] G. Furtos, M. Tomoaia-Cotisel, C. Garbo, M. Senila, N. Jumate, I. Vida-Simiti, C. Prejmerean, New composite bone cement based on hydroxyapatite and nanosilver, *Particulate Science and Technology*, **31**(4), 392-398 (2013). <https://doi.org/10.1080/02726351.2013.767293>
- [150] A. Mocanu, O. Cadar, P.T. Frangopol, I. Petean, Gh. Tomoaia, G.-A. Paltinean, C.P. Racz, O. Horovitz, M. Tomoaia-Cotisel, Ion release from hydroxyapatite and substituted hydroxyapatites in different immersion liquids: in vitro experiments and theoretical modelling study, *Royal Society Open Science*, **8**(1), 201785 (2021). <https://doi.org/10.1098/rsos.201785>
- [151] D. Oltean-Dan, G.-B. Dogaru, E.-M. Jianu, S. Riga, M. Tomoaia-Cotisel, A. Mocanu, L. Barbu-Tudoran, Gh. Tomoaia, Biomimetic composite coatings for activation of titanium implant surfaces: methodological approach and in vivo enhanced osseointegration, *Micromachines*, **12**(11), 1325 (2021). <https://doi.org/10.3390/mi12111352>
- [152] A. Mocanu, P.T. Frangopol, R. Balint, O. Cadar, I.M. Vancea, R. Mantiu, O. Horovitz, M. Tomoaia-Cotisel, Higuchi model applied to ions release from hydroxyapatites, *Studia UBB Chemia*, **66**(3), 195-207 (2021). <https://doi.org/10.24193/subbchem.2021.3.12>
- [153] A. Mocanu, G. Furtos, S. Rapuntean, O. Horovitz, C. Flore, C. Garbo, A. Danisteanu, Gh. Rapuntean, C. Prejmerean, M. Tomoaia-Cotisel, Synthesis characterization and antimicrobial effects of composites based on multi-substituted hydroxyapatite and silver nanoparticles, *Applied Surface Science*, **298**, 225-235 (2014). <https://doi.org/10.1016/j.apsusc.2014.01.166>
- [154] F. Goga, E. Forizs, G. Borodi, Gh. Tomoaia, A. Avram, R. Balint, A. Mocanu, O. Horovitz, M. Tomoaia-Cotisel, Behavior of doped hydroxyapatites during the heat treatment, *Revista de Chimie*, **68**(12), 2907-2913 (2017). <https://doi.org/10.37358/RC.17.12.6004>
- [155] P.T. Frangopol, A. Mocanu, V. Almasan, C. Garbo, R. Balint, G. Borodi, I. Bratu, O. Horovitz, M. Tomoaia-Cotisel, Synthesis and structural characterization of strontium substituted hydroxyapatites, *Revue Roumaine de Chimie*, **61**(4-5), 337-344 (2016).
- [156] A. Mocanu, R. Balint, C. Garbo, L. Timis, I. Petean, O. Horovitz, M. Tomoaia-Cotisel, Low crystallinity nanohydroxyapatite prepared at room temperature, *Studia UBB Chemia*, **62**(2), 95-103 (2017). <https://doi.org/10.24193/subbchem.2017.2.07>
- [157] R. Balint, I. Petean, P.T. Frangopol, A. Mocanu, G. Arghir, S. Riga, Gh. Tomoaia, O. Horovitz, M. Tomoaia-Cotisel, Biomimetic nanocomposite structures designed for coating of orthopedic implants: AFM investigation, *Studia UBB Chemia*, **66**(3), 141-160 (2021). <https://doi.org/10.24193/subbchem.2021.3.08>
- [158] F. Goga, E. Forizs, A. Rotaru, A. Lucian, I. Petean, A. Mocanu, M. Tomoaia-Cotisel, Synthesis and thermal treatment of hydroxyapatite doped with magnesium, zinc and silicon, *Revista de Chimie*, **68**(6), 1193-1200 (2017). <https://doi.org/10.37358/RC.17.6.5640>
- [159] C. Garbo, J. Locs, M. D'Este, G. Demazeau, A. Mocanu, C. Roman, O. Horovitz, M. Tomoaia-Cotisel, Advanced Mg, Zn, Sr, Si multi-substituted hydroxyapatites for bone

- regeneration, *International Journal of Nanomedicine*, **15**, 1037-1058 (2020). <https://doi.org/10.2147/IJN.S226630>
- [160] D. Oltean-Dan, G.B. Dogaru, M. Tomoaia-Cotisel, D. Apostu, A. Mester, H.R.C. Benea, M. G. Paiusan, E. M. Jianu, A. Mocanu, R. Balint, C.O. Popa, C. Berce, G.I. Bodizs, A.M. Toader, Gh. Tomoaia, Enhancement of bone consolidation using high-frequency pulsed electromagnetic short-waves and titanium implants coated with biomimetic composite embedded into PLA matrix: in vivo evaluation, *International Journal of Nanomedicine*, **14**, 5799-5816 (2019). <https://doi.org/10.2147/IJN.S205880>
- [161] A. Avram, M. Gorea, R. Balint, L. Timis, S. Jitaru, A. Mocanu, M. Tomoaia-Cotisel, Portland cement enriched with hydroxyapatite for endodontic applications, *Studia UBB Chemia*, **62**(4), 81-92 (2017). <https://doi.org/10.24193/subbchem.2017.4.07>
- [162] S. Rapuntean, P.T. Frangopol, I. Hodisan, Gh. Tomoaia, D. Oltean-Dan, A. Mocanu, C. Prejmerean, O. Soritau, L.Z. Racz, M. Tomoaia-Cotisel, In vitro response of human osteoblasts cultured on strontium substituted hydroxyapatites, *Revista de Chimie*, **69**(12), 3537-3544 (2018). <https://doi.org/10.37358/RC.18.12.6787>
- [163] C. Garbo, M. Sindilaru, A. Carlea, Gh. Tomoaia, V. Almasan, I. Petean, A. Mocanu, O. Horovitz, M. Tomoaia-Cotisel, Synthesis and structural characterization of novel porous zinc substituted nanohydroxyapatite powders, *Particulate Science and Technology*, **35**(1), 29-37 (2017). <https://doi.org/10.1080/02726351.2015.1121180>
- [164] A. Bogdanova, A. Makhro, J. Wang, P. Lipp, L. Kaestner, Calcium in red blood cells—a perilous balance, *International Journal of Molecular Sciences*, **14**(5), 9848-9872 (2013). <https://doi.org/10.3390%2Fijms14059848>
- [165] T.R.L. Klei, J.J. Dalimot, B.M. Beuger, M. Veldthuis, F.A. Ichou, P.J.J.H. Verkuijlen, I.M. Seignette, P.C. Ligthart, T.W. Kuijpers, R. Van Zwieten, R. Van Bruggen, The Gardos effect drives erythrocyte senescence and leads to Lu/BCAM and CD44 adhesion molecule activation, *Blood Advances*, **4**(24), 6218-6229 (2020). <https://doi.org/10.1182%2Fbloodadvances.2020003077>
- [166] L. Kuck, J.N. Peart, M.J. Simmonds, Calcium dynamically alters erythrocyte mechanical response to shear, *Biochimica et Biophysica Acta (BBA) – Molecular Cell Research*, **1867**(11), 118802 (2020). <https://doi.org/10.1016/j.bbamcr.2020.118802>
- [167] F. Jung, S. Braune, Thrombogenicity and hemocompatibility of biomaterials, *Biointerphases*, **11**(2), 029601 (2016). <https://doi.org/10.1116/1.4938557>
- [168] Y. Han, X. Wang, H. Dai, S. Li, Nanosize and surface charge effects of hydroxyapatite nanoparticles on red blood cell suspensions, *ACS Applied Materials & Interfaces*, **4**, 4616–4622 (2012). <https://doi.org/10.1021/am300992x>
- [169] T.P. Gamagedara, M.R.F. Riska, R.M.G. Rajapakse, Effects of hydroxyapatite - polymethyl methacrylate nanocomposites on human red cell indices and serum proteins, *International Journal of Nano Research*, **2**(1), 9-14 (2019).
- [170] C. Santos, S. Turiel, P. Sousa Gomes, E. Costa, A. Santos-Silva, P. Quadros, J. Duarte, S. Battistuzzo, M.E. Fernandes, Vascular biosafety of commercial hydroxyapatite particles: discrepancy between blood compatibility assays and endothelial cell behavior, *Journal of Nanobiotechnology*, **16**, 27 (2018). <https://doi.org/10.1186/s12951-018-0357-y>
- [171] N. Puvvada, P.K. Panigrahi, A. Pathak, Room temperature synthesis of highly hemocompatible hydroxyapatite, study of their physical properties and spectroscopic correlation of particle size, *Nanoscale*, **2**, 2631–2638 (2010). <https://doi.org/10.1039/c0nr00611d>
- [172] M.S. Laranjeira, A. Moco, J. Ferreira, S. Coimbra, E. Costa, A. Santos-Silva, P.J. Ferreira, F.J. Monteiro, Different hydroxyapatite magnetic nanoparticles for medical imaging: Its effects on hemostatic, hemolytic activity and cellular cytotoxicity, *Colloids and Surfaces B: Biointerfaces*, **146**, 363-374 (2016). <http://dx.doi.org/10.1016/j.colsurfb.2016.06.042>